



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 487/22, C07B 61/00, C07C 29/50, 31/12	A1	(11) International Publication Number: WO 96/17844 (43) International Publication Date: 13 June 1996 (13.06.96)
(21) International Application Number: PCT/US95/15899 (22) International Filing Date: 7 December 1995 (07.12.95) (30) Priority Data: 08/352,555 8 December 1994 (08.12.94) US (71) Applicant: THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; Center for Technology Transfer, Suite 300, 3700 Market Street, Philadelphia, PA 19104-3147 (US). (71)(72) Applicant and Inventor: DIMAGNO, Stephen [US/US]; 2828 Jackson Drive, Lincoln, NE 68502 (US). (72) Inventor: THERIEN, Michael, J.; 2411 D Delancey Place, Philadelphia, PA 19103 (US). (74) Agents: ELDERKIN, Dianne, B. et al.; Woodcock Washburn Kurtz MacKiewicz & Norris, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NOVEL PORPHYRINS AND PORPHYRIN SYNTHESIS TECHNIQUES (57) Abstract Strapped porphyrins and electron-deficient porphyrins are provided, as well as processes and intermediates for their preparation. In certain embodiments, such compounds are prepared by nucleophilic displacement of leaving groups from methylpyrroles. In other embodiments, such compounds are prepared by condensing pyrrole derivatives and removing water thus formed from the resulting reaction mixture.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

NOVEL PORPHYRINS AND PORPHYRIN SYNTHESIS TECHNIQUES

RELATED APPLICATIONS

This patent application is a continuation-in-part of the United States Patent Application entitled "Electron-
5 Deficient Porphyrins And Processes And Intermediates For Preparing Same," filed on April 28, 1994, which is a continuation-in-part of United States Application Serial No. 08/064,468, filed May 20, 1993, which, in turn, is a continuation-in-part of United States Application Serial No.
10 07/929,943, filed August 14, 1992. The entire contents of these patent applications are incorporated herein by reference.

FIELD OF THE INVENTION

This invention relates to novel chiral and/or
15 strapped porphyrins and to techniques and intermediates useful in preparing these and other types of porphyrin-based compounds.

BACKGROUND OF THE INVENTION

Porphyrins are derivatives of porphine, a conjugated
20 cyclic structure of four pyrrole rings linked through their 2- and 5-positions by methine bridges. Porphyrins can be covalently attached to other molecules. The electronic features of the porphyrin ring system can be altered by the attachment of one or more substituents. The term "porphyrin"
25 includes derivatives wherein a metal atom is inserted into the ring system, as well as molecular systems in which ligands are

- 2 -

attached to the metal. The substituents, as well as the overall porphyrin structure, can be neutral, positively charged, or negatively charged.

Electron-deficient porphyrins (i.e., porphyrins bearing substituents that are electron-withdrawing relative to hydrogen) have been suggested for use as industrial oxidation and reduction catalysts. Strapped porphyrins, wherein atoms at or near the periphery of the porphyrin ring system are covalently bound through linkages (straps) that do not include the porphyrin core, also have been suggested for a variety of uses, including use as enantioselective oxidation catalysts. A number electron-deficient porphyrins and strapped porphyrins have been prepared, but many known synthetic methods generally proceed in low yield, if at all, and cannot be used to produce many types of porphyrins.

Accordingly, there exists a need in the art for efficient synthetic methods capable of producing a greater variety of electron-deficient and/or strapped porphyrins.

OBJECTS OF THE INVENTION

It is one object of the present invention to provide improved methods for synthesizing electron-deficient porphyrins.

It is another object of the invention to provide improved methods for synthesizing strapped and/or chiral porphyrins.

It is yet another object to provide novel electron-deficient porphyrins.


It is yet another object to provide novel strapped porphyrins.

It is still another object to provide synthetic precursors for electron-deficient, strapped and/or chiral porphyrins.


SUMMARY OF THE INVENTION

These and other objects are satisfied by the present invention, which provides novel processes for preparing

porphyrins and novel strapped and/or electron-deficient porphyrins prepared by such processes. In certain embodiments, porphyrins are prepared by contacting a pyrrole derivative having formula (1) with a pyrrole compound having
 5 formula (2) in organic solvent for a time and under conditions effective to form a covalent linkage between the pyrrole derivative and the pyrrole.



(1)




(2)

In preferred embodiments R_A and R_B are selected from the group consisting of H, groups that are electron-withdrawing relative
 10 to hydrogen, covalently-bound multi-atom linkages, and acid-stable chemical functional groups such that: at least one R_A is H or a group that is electron-withdrawing relative to hydrogen; at least two R_A , together, form a covalently-bound, multi-atom linkages; at least two R_B , together, form a
 15 covalently-bound, multi-atom linkage; or at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage. In preferred embodiments, R_A has formula CF_2R , wherein R is alkyl, alkenyl, alkynyl, or a group having a higher formal oxidation state, such as a halogenated or
 20 perhalogenated derivative. In preferred embodiments, the multi-atom linkage includes a group that is electron-withdrawing relative to hydrogen. L is a leaving group and n preferably is 0, 1, or 2.


In further embodiments, porphyrins are prepared by
 25 reacting at least two molecules having formula (3) in organic solvent for a time and under conditions effective to form a covalent bond between said molecules.

- 4 -




(3)

The invention also provides synthetic methods wherein an aldehyde having formula $R_A\text{-CHO}$ is contacted with a pyrrole derivative having formula (4) ($q = 0, 1, \text{ or } 2$) in organic solvent in the presence of acid for a time and under conditions effective to form a reaction mixture comprising water and an adduct of the aldehyde and the pyrrole derivative, and a portion of the water thus formed is removed from said reaction mixture.



(4)

Also provided are synthetic methods wherein an alcohol having formula (5) is reacted in organic solvent in the presence of acid for a time and under conditions effective to form a reaction mixture comprising water and an adduct of the alcohol, and a portion of the water thus formed is removed from the reaction mixture.




(5)

In other embodiments, porphyrins are prepared by contacting a pyrrole derivative having formula (6) with a


- 5 -

pyrrole compound having formula (2) in organic solvent in the presence of acid for a time and under conditions effective to form a reaction mixture comprising water and an adduct of the pyrrole derivative and then removing from the reaction mixture
 5 a portion of the water thus formed.




(6)


The invention also provides strapped porphyrins having formula (7), (8), or (9):



(7)



(8)



(9)

wherein M and M' are metal atoms and at least two R_A, at least two R_B, or an R_A and an R_B, together, form a group that is

- 6 -

electron-withdrawing relative to hydrogen. In preferred embodiments, at least two R_A and/or at least two R_B are alkyl having about 3 to about 35 carbon atoms, aryl having about 6 to about 40 carbon atoms, $C(O)O-(alkyl)$ or $C(O)-(alkyl)$.

5 In another aspect, the invention provides polymers comprising linked porphyrin units, at least one of such units being a strapped porphyrin. In certain embodiments, porphyrin units having formula (7), (8), or (9) share covalent bonds. In other embodiments, at least one R_A group or R_B group
10 functions as a linking group. In these embodiments, at least a portion of a linking group can have formula $[C(R_C)=C(R_D)(R_E)]_x$, $[C\equiv C(R_D)]_x$, $[CH_2(R_C)-CH(R_D)(R_E)]_x$ or $[CH=CH(R_D)]_x$ where R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon
15 atoms, aryl or heteroaryl having about 4 to about 20 carbon atoms, alkenyl or heteroalkenyl having from 1 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 1 to about 20 carbon atoms, trialkylsilyl or porphyrinato, provided that at least one of R_C , R_D , and R_E is not H and x is at least 1.
20 R_C , R_D , and R_E also can include peptides, nucleosides, and/or saccharides. The remaining of R_A and R_B can be H, halogen, alkyl or heteroalkyl having 1 to about 20 carbon atoms or aryl or heteroaryl having 4 to about 20 carbon atoms, $C(R_C)=C(R_D)(R_E)$, $C\equiv C(R_D)$, or a chemical functional group that
25 includes a peptide, nucleoside, and/or saccharide. In other preferred embodiments, the linking group is cycloalkyl or aryl having about 6 to about 22 carbon atoms.

The invention also provides processes for preparing porphyrin-containing polymers. In certain embodiments, the
30 processes comprise providing at least two compounds that, independently, have formula (7), (8), or (9) wherein at least one R_A group or R_B group in each of the compounds contains an olefinic carbon-carbon double bond or a chemical functional group reactive therewith. In other embodiments, at least one
35 R_A group or R_B group in each of the compounds contains a carbon-carbon triple bond or a chemical functional group reactive therewith. The compounds are then contacted for a

-7-

time and under reaction conditions effective to form covalent bonds through the carbon-carbon double and/or triple bonds.

The porphyrins and porphyrin-containing polymers of the invention can be used, for example, as dyes, catalysts, contrast agents, antitumor agents, antiviral agents, and in chemical sensors and electrooptical devices.

BRIEF DESCRIPTION OF THE DRAWINGS

The numerous objects and advantages of the present invention may be better understood by those skilled in the art by reference to the accompanying figures, in which:

Figure 1 shows a synthetic scheme for the preparation of chiral tetraarylporphyrins.

Figure 2 shows a synthetic scheme for the preparation of doubly-strapped porphyrins.

DETAILED DESCRIPTION OF THE INVENTION

It has been found in accordance with the present invention that a wide variety of strapped porphyrins can be prepared by condensation of suitably functionalized pyrroles provided that at least a portion of the water of condensation is removed from the reaction mixture. In general, the resulting porphyrins have formula (7), (8), or (9) wherein M and M' are metal atoms and R_A and/or R_B groups, together, form a covalently-bound, multi-atom linkage. The remaining R_A and R_B groups can be the same or different and are selected from H and those groups known to be stable under the reaction conditions (e.g., acidic reaction conditions) under which such compounds are formed, including alkyl, alkenyl, alkynyl, and aryl groups. Representative groups are disclosed by, for example, Application Serial No. 08/064,468. Those skilled in the art will recognize that chemical protecting groups can be attached to sensitive functionality found within R_A and/or R_B and that such protecting groups can be removed after a particular reaction has been completed. (see, e.g., Greene and Wuts in Protective Groups in Organic Synthesis, New York, John Wiley & Sons, 1991).

- 8 -

Compounds having formulas (7)-(9) preferably bear 1, 2, 4, 8, or 12 substituents (i.e., 1, 2, 4, 8, or 12 of R_A and R_B are not H). In certain embodiments, four R_A groups bear electron-withdrawing functionality. In other
5 embodiments, a single R_B on each pyrrol moiety is a halogen or both R_B on two non-adjacent pyrrol moieties are halogens.

Numerous examples of electron-withdrawing functional groups are known to those skilled in the art. Further, electron-withdrawing groups can be identified through routine
10 experimentation involving, for example, replacement of hydrogen in a molecule with a given group and then testing any resultant inductive effects. Representative electron withdrawing groups include the following: $N-(alkyl)_3$, NH_3^+ , NO_2 , $SO_2-(alkyl)$, CN , $SO_2-(aryl)$, $C(O)OH$, F , Cl , Br , I , CF_2R_1 ,
15 CCl_2R_1 , CBr_2R_1 , CI_2R_1 , $C(O)O-(alkyl)$, $C(O)-(alkyl)$, and/or CHO , wherein alkyl groups have from about 1-35 carbon atoms and aryl groups have about 3-50 carbon atoms. Electron-withdrawing groups used as non-terminal substituents (e.g., those substituents that form strapping groups) may preferably
20 include at least one chiral center. Alkyl and aryl groups found in non-terminal substituents have from about 1 to about 50 carbon atoms and about 3 to about 75 carbon atoms, respectively. Alkenyl and alkynyl groups found in non-terminal substituents have from about 2 to about 50 carbon
25 atoms. Preferably, such alkyl, alkenyl, and alkynyl groups have from 3 to about 35 carbon atoms and such aryl groups have from about 6 to about 40 carbon atoms. More preferably, such alkyl, alkenyl, and alkynyl groups have from 6 to about 24 carbon atoms and such aryl groups have from about 8 to about
30 30 carbon atoms. Alkyl and aryl groups found in terminal substituents preferably have from 1 to about 20 carbon atoms and about 6 to about 20 carbon atoms, respectively. Alkenyl and alkynyl groups found in terminal substituents have from about 2 to about 20 carbon atoms. More preferably, such
35 alkyl, alkenyl, and alkynyl groups have from 5 to about 20 carbon atoms and such aryl groups have from about 6 to about 20 carbon atoms. The terms alkyl, alkenyl, alkynyl, and aryl

- 9 -

are intended to include moieties substituted with, for example, halogens or nitro groups, as well as moieties wherein heteroatoms (e.g., N, O, S, Se, and Te) are inserted into the carbon backbone of an alkyl or aryl structure to yield, for example, an ether, thioether, and pyridinyl group. Alkyl, alkenyl, alkynyl, and aryl groups can bear substituents that include additional carbon atoms. Preferred electron-withdrawing groups are substituted and unsubstituted alkyl and aryl groups that possess net electron-withdrawing effects.

10 Perhaloalkyl and perhaloaryl groups are particularly preferred, including perfluoroalkyl, perfluorophenyl, perfluorobenzyl, and tetrafluoropyridyl groups.

Numerous acid stable groups are known to those skilled in the art. (see, e.g., March, Advanced Organic Chemistry, 4th ed., 1992). Representative acid stable groups include halogens, NO₂, and CN.



15

M preferably is a lanthanide or actinide or a metal such as Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Y, Zr, Nb, Mo, Ru, Re, Rh, Pd, Ag, La, Hf, Ta, W, Os, Ir, Pt, Cd, Hg, Li or Au. More preferably, M is Cr, Mn, Fe, Co, Ni, Ru, Re, Rh, Pd, Os, Ir, Pt, or Au. M' can be a metal such as Li, Na, K, Rb, or Cs, preferably Li.

20


In certain embodiments, porphyrins according to the invention are prepared by synthesizing and then oxidizing suitably-substituted porphyrinogen compounds having, for example, formulas (10) and (11).

25




-10-


In other embodiments porphyrins are prepared by condensation and subsequent oxidation of suitably-substituted polypyrrolyl intermediates having, for example, formulas (12)-(15).




(12)



(13)




(14)




(15)

5 Porphyrinogens and polypyrrolyl intermediates can be prepared by reacting pyrrole derivatives having formula (1) with pyrrole compounds having formula (2), or by reacting at least two molecules of pyrrole derivatives having formula (3).


-11-



(1)



(2)



(3)


These reactions should be performed in organic solvent for a time and under conditions effective to form a covalent linkage between an unsubstituted 2- or 5-position in a first pyrrole compound (e.g., formula (2)) and an exocyclic methylene carbon atom in a second pyrrole compound (e.g., formula (1)). While not wishing to be bound by any particular theory, it is believed that such reactions proceed by loss of leaving group, L, from the exocyclic methylene by an S_N1 process. Thus, preferred L groups facilitate such reactions by, for example, readily forming stable (and, preferably, inert) anionic species. A wide variety of leaving groups can be used in accordance with the present invention, including those disclosed by Advanced Organic Chemistry, J. March, 4th ed., John Wiley & Sons, New York, 1992. Some leaving groups may need to be protonated before displacement. Preferred leaving groups include halogens, sulfates, sulfonates (e.g., methanesulfonate, trifluoromethanesulfonate, toluenesulfonate, and *p*-bromobenzenesulfonate), carboxylates (e.g., acetate, trifluoroacetate, and *p*-nitrobenzoate), and silyl ethers (e.g., trimethylsilylethers).

Condensation reactions according to the invention can be performed in a wide variety of organic solvents, including trichloroethylene, chloroform, carbon tetrachloride,


-12-

benzene, toluene, xylenes, methylene chloride, dimethylsulfoxide, acetonitrile, dimethylformamide, and anisole. Aprotic solvents are preferred. In preferred embodiments, such reactions are performed between 0°C and 130°C.

Alternatively, porphyrinogens and polypyrryl intermediates can be prepared by condensing alcohols having formula (5) ($n = 0, 1, \text{ or } 2$). Such compounds also can be prepared by condensing aldehydes having formula $R_A\text{-CHO}$ with a pyrrole derivative having formula (4) ($q = 0, 1, \text{ or } 2$).




(4)



(5)

Porphyrinogens and polypyrryl intermediates alternatively can be prepared by condensing alcohols having formula (6) ($n = 0, 1, \text{ or } 2$) with pyrrole compounds having formula (2).



(6)

Each of these condensation reactions should be performed in organic solvent in the presence of acid for a time and under conditions effective to form reaction mixtures comprising water and an adduct of the reagents. In accordance with the invention, at least a portion of the water thus formed is removed from the reaction mixture.

A wide variety of organic solvents can be used in the condensation reactions of the invention, including benzene, toluene, xylenes, methylene chloride, chloroform, trichloroethylene, and mixtures thereof. Aprotic solvents are

-13-

preferred, particularly nonpolar, aprotic solvents. Solvents capable of forming azeotropes (i.e., constant boiling mixtures) with water are particularly preferred. In preferred embodiments, condensation reactions are performed between 35°C and 110°C.

Acids according to the invention are ions or molecules having the capacity to accept at least one electron pair. Representative acids include benzoic acid, sulfonic acids (e.g., *p*-toluenesulfonic acid and methanesulfonic acid), trifluoroacetic acid, boron trifluoride, boron trichloride, and mixtures thereof. Preferred acids are not volatile under reactions conditions of the invention. Protic acids, particularly strong protic acids (i.e., those having $pK_a < 0$), are preferred. In preferred embodiments, a catalytic (i.e., non-stoichiometric) amount of acid is used.

Water can be removed from adduct-containing reaction mixtures by a wide variety of known techniques, including membrane-based separations. Water also can be removed by contacting a reaction mixture with moieties that absorb, trap, or react with water or otherwise render water non-reactive. In general, the chosen technique should remove at least a portion of any water present but should not remove the adduct-forming reagents. Representative water removal techniques are disclosed by U.S. Patent No. 4,332,643 (Reid), European Patent Application EP 92-114390 (Inaba, et al.), Japanese Patent Applications 91-146674 (Miyazaki, et al.), 91-20083 (Kondo, et al.), and 90-104128 (Okazaki, et al.), and Brazilian Patent Application 77-433 (Scaglia, et al.). Water preferably is removed by distilling an azeotrope formed by the water and the organic solvent. In certain embodiments, the distilled azeotrope is collected in a vessel and allowed to separate into aqueous and organic phases, and the organic (solvent) phase is returned to the reaction mixture. In other embodiments, the distilled azeotrope is contacted with a drying agent and the dried distillate is returned to the reaction mixture. Representative drying agents include phosphorous pentoxide, calcium hydride, calcium oxide, barium

-14-

oxide, lithium aluminum hydride, molecular sieves, and mixtures thereof. Numerous additional drying agents are well-known to persons of ordinary skill in the art. In further embodiments, the dried distillate is collected and a roughly
5 equal volume of fresh solvent is added to the reaction mixture. In still further embodiments, semi-permeable membrane technology is used to remove water from the reaction mixture as it is formed.

Compounds having formulas (4)-(6) preferably
10 prepared by contacting a pyrrole having formula (2) with base with or without organic solvent in the presence of aldehydes having formula $R_A\text{-CHO}$. Representative bases include sodium hydroxide, lithium hydroxide, potassium hydroxide, barium hydroxide, alkyl or aryl lithium reagents, and alkyl or aryl
15 Grignard reagents, with sodium hydroxide being preferred. Representative solvents include tetrahydrofuran, ethers, alcohols, water, dimethylsulfoxide, and dimethylformamide. In preferred embodiments, such reactions are performed between -78°C and 30°C . The pyrrole, aldehyde, and base can be
20 reacted simultaneously or in a number of different ways. For example, the pyrrole can be contacted with base and then added to the aldehyde or it can be contacted with base in the presence of aldehyde. In certain embodiments, pyrrole, aldehyde, and base are contacted in the absence of solvent.
25 The product produced depends, in part, upon the proportions of pyrrole and aldehyde used. For example, the use of excess pyrrole tends to favor formation of compound (4), the use of excess aldehyde tends to favor formation of compound (6), and the use of equimolar proportions tends to favor formation of
30 compound (5).

Compounds having formulas (1) and (2) preferably are prepared by reacting hydroxymethylpyrroles having formulas (5) and (6) with suitable derivatizing reagents, preferably in organic solvent and in the presence of base. Representative
35 bases include trimethylamine, triethylamine, pyridine, dimethylaminopyridine, and 1,8-bis(dimethylamino)naphthalene, N,N,N',N'-tetramethyl-1,8-naphthalenediamine(Proton-Sponge[®],

-15-

Aldrich Chemical Co.). Representative solvents include methylene chloride, chloroform, and tetrahydrofuran. In preferred embodiments, such reactions are performed between 0°C and 25°C. A wide variety of useful derivatizing reagents are known in the art. Preferred derivatizing reagents react with the hydroxyl oxygen atom to produce leaving group L. Representative of such derivatizing reagents are the sulfonyl halides, silyl halides, acid halides, and acid anhydrides.

Oxidation of porphyrinogens and polypyrryl intermediates can be accomplished by a number of techniques. For example, porphyrinogen- and/or polypyrryl-containing reaction mixtures can be exposed to oxidizing conditions. Alternatively, such compounds are isolated from a reaction mixture and then contacted with an oxidizing agent. Representative oxidizing agents include oxygen, *p*-chloranil, 2,3-dichloro-5,6-dicyanobenzoquinone (DDQ), and mixtures. Oxidation of electron-deficient compounds also can be effected using bulk electrochemical methods (see, e.g., Laboratory Techniques in Electroanalytical Chemistry, P.T. Kissinger and W.R. Heineman, eds., New York, Marcel Dekker, 1984). In general, oxidation conditions for partially-oxidized porphyrinogens and polypyrryl intermediates (e.g., formulas (11), (13), and (15)) will be less vigorous than for porphyrinogens and polypyrryl intermediates in more reduced form (e.g., formulas (10), (12), and (14)). More electron-deficient porphyrinogens generally require more vigorous oxidation conditions. In preferred embodiments, oxidation reactions are performed between 25°C and 150°C.

The processes of the invention produce somewhat monomeric compounds that can be incorporated into porphyrin-containing homopolymers or copolymers or into macromolecular or supramolecular species containing, for example, one or more peptides, nucleosides, or saccharides. Polymers according to the invention can contain as few as 2 porphyrin units, but more preferably contain at least 3 porphyrin units, more preferably at least 5 porphyrin units. In certain embodiments, polymers of the invention comprise a plurality


-16-

of porphyrin units that, independently, have formula (7), (8), or (9) wherein at least one R_A group or R_B group includes a linking group selected from $[C(R_C)=C(R_D)(R_E)]_x$, $[C\equiv C(R_D)]_x$, $[CH_2(R_C)-CH(R_D)(R_E)]_x$ or $[CH=CH(R_D)]_x$ where x is at least 1.

- 5 The remaining R_A and R_B include at least one group that is electron-withdrawing relative to hydrogen.

In other embodiments, polymers according to the invention comprise a plurality of porphyrin units that, independently, have formula (7), (8), or (9) wherein at least
10 one R_A group or R_B group is a cycloalkyl, cycloalkenyl, aryl or heteroaryl linking group having about 6 to about 22 carbon atoms.

Those skilled in the art will recognize the wide variety of polymers that can be prepared from the porphyrin-
15 containing compounds of the invention. In certain embodiments, cofacial polymers are formed having, for example, formula (16). (see, e.g., Durand, et al., *J. Am. Chem. Soc.*, 1983, 105, 2710).




(16)

20 In other embodiments, somewhat linear polymer chains are formed wherein a portion of the polymer has general formula $(P_N)_r$ where P_N is a porphyrin unit and r is at least 2. In further embodiments, linear polymer chains have general formula:


25
$$[-(Q_L)_t-(P_N)_s-]_h$$

where Q_L is a linking group, P_N is a porphyrin unit, and h , t , and s are independently selected to be at least 1. For example, a portion of such polymers can have formula:


-17-



wherein P_{N1} and P_{N2} are independently selected porphyrin units, Q_{L1} and Q_{L2} are independently selected linking groups, and l' , l'' , s' , and s'' are at least 1. These essentially linear
 5 polymer chains can be cross-linked such that a portion of the polymer has general formula:



wherein Q_H is a linking group, and h , u , and v are independently selected to be at least 1. A portion of these
 10 cross-linked polymers can have formula:



wherein P_{N3} is a porphyrin unit, Q_{H1} and Q_{H2} are independently selected linking groups, and h' , h'' , u' , and u'' are at least 1. Thus, cross-linked polymers can have formulas:



where r' is at least 1.

The polymers of the invention can be formed by
 25 contacting a substituted porphyrin with a second compound containing functionality that is reactive with the functionality contained within the porphyrin. Preferably, the porphyrin contains an olefinic carbon-carbon double bond, a carbon-carbon triple bond or some other reactive
 30 functionality. The contacting should be performed under conditions effective to form a covalent bond between the respective reactive functionalities. Preferably, porphyrin-containing polymers are formed by metal-mediated cross-coupling of, for example, dibrominated porphyrin units. Also,
 35 porphyrin-containing polymers can be synthesized using known terminal alkyne coupling chemistry. (see, e.g., Patai, et al., The Chemistry of Functional Groups, Supplement C, Part 1, pp. 529-534, Wiley, 1983; Cadot, et al., Acetylenes, pp. 597-647, Marcel Dekker, 1964; and Eglinton, et al., Adv. Org. Chem.,

-18-

1963, 4, 225) As will be recognized, the second compound noted above can be a substituted porphyrin of the invention or some other moiety such as an acrylate monomer. Thus, a wide variety of copolymeric structures can be synthesized with the porphyrins of the invention. Through careful substituent selection the porphyrins of the invention can be incorporated into virtually any polymeric matrix known in the art, including but not limited to polyacetylenes, polyacrylates, polyolefins, polyethers, polyurethanes, polycarbonates, polyanilines, polypyrroles, and polythiophenes. For example, fluorescent porphyrins can be incorporated into such polymers as end-capping groups.

The porphyrins and porphyrin-containing polymers of the invention can be used, for example, as dyes, catalysts, contrast agents, antitumor agents, antiviral agents, liquid crystals, in chemical sensors and in electrooptical and solar energy conversion devices. One preferred use for compounds containing electron-deficient porphyrins are as catalysts for the oxygenation of alkanes and/or alkenes, particularly oxygenations performed in supercritical carbon dioxide. Electron-deficient porphyrins also can be incorporated into supramolecular structures. The polymers and supramolecular structures, which anchor porphyrin units in a relatively stable geometry, should improve many of the known uses for porphyrins and even provide a number of new uses, such as in a solid phase system for sterilizing virus-containing solutions. Representative uses are disclosed by, for example, the following patents, which are incorporated herein by reference: U.S. Patent No. 4,895,682 (Ellis, et al.); U.S. Patent No. 4,986,256 (Cohen); U.S. Patent No. 4,668,670 (Rideout, et al.); U.S. Patent No. 3,897,255 (Erickson); U.S. Patent No. 3,899,334 (Erickson); U.S. Patent No. 3,687,863 (Wacher); U.S. Patent No. 4,647,478 (Formanek, et al.); and U.S. Patent No. 4,957,615 (Ushizawa, et al.). Further uses are disclosed are disclosed by, for example, U.K. Patent Application 2,225,963 (Casson, et al.); International Application WO 89/11277 (Dixon, et al.); International

-19-

Application WO 91/09631 (Matthews, et al.); European Patent Application 85105490.8 (Weishaupt, et al.); European Patent Application 90202953.7 (Terrell, et al.); European Patent Application 89304234.1 (Matsushima, et al.); Lehn, *Angew. Chem. Int. Ed. Engl.*, 1988, 27, 89; Wasielewski, *Chem. Rev.*, 1992, 92, 435; Mansury, et al., *J. Chem. Soc., Chem. Comm.*, 1985, 155; Groves, et al., *J. Am. Chem. Soc.*, 1983, 105, 5791; and Giroud-Godquin, et al., *Angew. Chem. Int. Ed. Engl.*, 1991, 30, 375. It is believed that the porphyrins of the invention can be substituted for the porphyrins disclosed in each of the foregoing publications.

Additional objects, advantages, and novel features of this invention will become apparent to those skilled in the art upon examination of the following examples thereof, which are not intended to be limiting. Unless otherwise stated, all reactions are performed using dry solvents under inert atmospheres. Porphyrin-forming condensation reactions are all assumed to be carried out using continuous water removal methods.

Example 1

Preparation Of 2-(2',2',3',3',4',4',4'-Heptafluoro-1'-hydroxybutyl)pyrrole From Heptafluorobutyraldehyde Hydrate

Heptafluorobutyraldehyde hydrate (9.26 g, 42.9 mmol) was placed in a 100 mL Schlenk flask. This was frozen with liquid nitrogen and an inert atmosphere was established. Against an outflow of nitrogen, dried pyrrole (5.95 mL, 85.8 mmol) and sodium hydroxide (4.52 g, 113 mmol) were added. The flask was wrapped in foil and the mixture was stirred overnight, during which time it solidified. A colorless liquid also was present. The volatiles were removed by vacuum, leaving a light brown solid. The solid was dissolved in 40 mL of water and the solution was extracted (4 x 50 mL) with methylene chloride. The organic layers were dried over sodium sulfate and then evaporated to dryness under vacuum to give 5.39 g (47%) of a light yellow-brown solid. ¹H NMR,

-20-

(CDCl₃, 360 MHz) d 8.50 br 1H; 6.87 m 1H; 6.32 m 1H; 6.22 m 1H; 5.28 d, J = 8.17 Hz; 5.23 d, J = 7.67 Hz; 2.42 br s.

Example 2

Preparation Of 2-(2',2',3',3',4',4',4'-Heptafluoro-1'-hydroxybutyl)pyrrole From Using Organolithium Reagents

Dry, distilled pyrrole (80 mmol) is dissolved in diethyl ether (200 ml) and cooled to -78°C. Butyl lithium (80 mmol, 32 ml of 2.5 M solution in hexane) is added dropwise with stirring and the solution is gradually warmed to room temperature with evolution of hydrogen. This solution is transferred dropwise by cannula to a -78°C solution of dry heptafluorobutyraldehyde (previously distilled from P₂O₅) in tetrahydrofuran (THF). The solution is warmed to room temperature with stirring. The volatiles are removed by vacuum leaving a solid that is dissolved in 40 mL of water and extracted (4 x 50 ml) with methylene chloride. The organic layers are dried over sodium sulfate and then evaporated to dryness under vacuum to give the product.

Example 3

Preparation Of

2,5-Bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole

Heptafluorobutyraldehyde hydrate (5.0 g, 23 mmol) was placed in a 100 mL Schlenk flask. This was frozen with liquid nitrogen and an inert atmosphere was established. Against an outflow of nitrogen, dried pyrrole (0.694 mL, 10 mmol) and sodium hydroxide (2.25 g, 56 mmol) were added. The flask was wrapped in foil and the mixture was stirred for 2 days. The volatiles were removed by vacuum, leaving a light brown, oily solid. The solid was dissolved in 40 mL of water and the solution was extracted (4 x 50 ml) with methylene chloride. The organic layers were dried over sodium sulfate and then evaporated to dryness under vacuum to an oily brown solid (38%), which proved to be a diastereomeric mixture of the desired products. 1H NMR of recrystallized product (CDCl₃,

-21-

360 MHz) d 8.78 (broad s, 1 H); 6.28 (d, 2 H, J = 2.8 Hz); 5.128 (dd, 2 H, J1 = 117 Hz, J2 = 8 Hz); 2.53 (broad s, 2 H).

Example 4**Preparation Of****5 2-(2',2'-Difluoro-2'-pentafluorophenyl-1'-hydroxyethyl)pyrrole**

2-Pentafluorophenyl-2,2-difluoroethanal (40 mmol) and THF (5 ml) are placed in a 100 mL Schlenk flask. This is frozen with liquid nitrogen and an inert atmosphere is established. Against an outflow of nitrogen, dried pyrrole
10 (5.95 mL, 85.8 mmol) and sodium hydroxide (4.52 g, 113 mmol) is added. The flask is wrapped in foil and the mixture is stirred overnight. The volatiles are removed by vacuum, the resulting solid is dissolved in 40 mL of water, and the solution is extracted (4 x 50 ml) with methylene chloride.
15 The organic layers are dried over sodium sulfate and then evaporated to dryness under vacuum to give 2-(2,2-difluoro-2-pentafluorophenyl-1-hydroxyethyl)pyrrole.

Example 5**Preparation Of 2-(2',2',2'-Trifluoro-1'-hydroxyethyl)pyrrole**

20 The procedure of Example 4 is repeated except that trifluoroacetaldehyde is used in place of 2-Pentafluorophenyl-2,2-difluoroethanal.

Example 6**Preparation Of 1-Pyrrolyl-1-perfluoroundecyl Methanol**

25 The procedure of Example 4 is repeated except that perfluorododecanal is used in place of 2-Pentafluorophenyl-2,2-difluoroethanal.

Example 7**Preparation Of Tetrakis(heptafluoropropyl)porphyrin From 1-Pyrrolyl-1-perfluoropropyl-Methanol**

30 Benzene (650 ml) was placed in a one liter, double-necked flask and azeotropically dried under nitrogen using a recycling Dean-Stark apparatus. p-Toluenesulfonic acid

-22-

hydrate (50 mg) was added to the benzene and azeotropic distillation was continued until the distillate stopped phase separating. The Dean-Stark trap was emptied and 4Å molecular sieves (20 ml) were added to the trap. Distillation was continued for 10 minutes with the distillate recycling through the molecular sieves. 2-Pyrrolylperfluoropropylmethanol (265 mg, 1 mmol) was dissolved in 10 ml of dry benzene and added (all at once) to the benzene solution heated at reflux. The solution became pink immediately after the addition, then gradually darkened. Heating was continued for 30 minutes and the reaction mixture was quenched with 600 mg of DDQ. Heating was continued for an additional hour under N₂. The solution was transferred to a 1 liter round bottom flask and the solvent was removed and recovered by rotary evaporation. The remaining dark brown residue was dissolved, to the extent possible, in 50 ml of warm hexane containing 1 ml of pyridine, and was poured directly on to a short (2 x 10 cm) column consisting of silica that was packed in hexane and topped with a 2 cm pad of Celite. Elution of the porphyrin was carried out with hexane. Collection was continued until the eluant became nearly colorless. The solvent was removed from the collected fraction and the resulting solid was washed with cold hexane (10 ml) and filtered to yield 90 mg (37%) of nearly pure 5,10,15,20-tetrakis(perfluoropropyl)porphyrin. An analytical sample was recrystallized from chloroform (-20°C) to yield crystals suitable for X-ray diffraction. ¹H NMR (360 MHz, CDCl₃) δ 9.50 (s, 8 H); -2.30 (s, 2 H). ¹⁹F NMR (DCD₃, CF₃COOH ext. std) δ -79.7 (t, 3 F); -80.9 (broad s, 2 F); -118.8 (broad s, 2 F). The ¹⁹F spectrum shows evidence of exchange behavior. The signal at -118.8 ppm sharpens to a broadened triplet upon warming the solution to 55°C. ¹³C NMR (75 MHz, CDCl₃) gave only two discernable signals at 144.2 and 133.8 after a 16 hour run.

-23-

Example 8**Preparation Of Tetrakis(heptafluoropropyl)porphyrin From Pyrrole And Heptafluorobutyraldehyde**

In a procedure analogous to that described in Example 7, the apparatus was charged with benzene (650 ml), p-toluenesulfonic acid hydrate (50 mg) and heptafluorobutyraldehyde hydrate (0.22g, 1 mmol). After refluxing the mixture for 1 hour, dry pyrrole (70 μ l, 1 mmol) was added. The reaction was monitored by thin layer chromatography (TLC); after 1.5 hours the reaction was quenched as in Example 7. The reaction mixture was neutralized with pyridine, filtered through silica gel, pumped dry, and further purified by chromatography on silica. Several pyrrole-containing products can be isolated from this preparation. The desired product, tetrakis(heptafluoro)-porphyrin, eluted as the first colored band. This method gives 4 mg (1.6%) of the target porphyrin.

Example 9**Preparation Of Tetrachloroporphine**

Porphine (Zn) (40 mg) was dissolved in 300 mL of a 1:1 mixture of THF and CHCl_3 , and the mixture was placed in a 500 ml round bottomed flask. N-Chlorosuccinimide (NCS) was added (4.2 eq.) and the mixture was stirred overnight protected from the light. The reaction was monitored by TLC and four intermediates were observed, presumably the target compound and the mono-, di-, and trichloro intermediates. After 24 hours the reaction was stopped and tetrachloroporphine (> 80%) was isolated.

Example 10**Preparation Of Cofacial Porphyrin Dimers**

To a THF solution of 5-bromo-10,15,20-trichloroporphyrinate (Zn) (1 eq.) is added $\text{Pd}(\text{P}(\text{C}_6\text{H}_5)_3)_2$ (5 mol%) and anthracene-1,8-bis(chlorozinc) (0.5 eq). The reaction is stirred for 24 hours at room temperature. One band is evident by TLC of the

-24-

reaction mixture. The compound is purified by silica gel chromatography to isolate the dimeric, anthracene bridged compound.

Example 11

5 Preparation Of 1,8-(2',5'-pyrrolyl)-1,8-hydroxy-2,2,3,3,4,4,5,5,6,6-dodecafluorooctane

Pyrrole (4.0 mmol) and 2,2,3,3,4,4,5,5,6,6,7,7-dodecafluoro-1,8-octanedicarboxaldehyde (20 mmol) are dissolved in 50 ml of THF. Sodium hydroxide (120 mmol) is added as pellets. The mixture is stirred at room temperature for one week. The solvent is removed in vacuo and the remaining solid is extracted from water with 3 x 50 ml of methylene chloride. The organic layer is dried over sodium sulfate and filtered, and the solvent is removed in vacuo, leaving the α,α' -perfluoroalkyl-strapped pyrrole and some polymeric side products. The mixture is purified on silica using hexane/diethyl ether as eluent. The product is isolated and the solvent is removed to leave the desired strapped pyrrole.

20 Example 12

Preparation of 2,5-Bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsiloxybutyl)pyrrole

Recrystallized 2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole (0.5 mmol, 0.2316 g) was dissolved in 25 ml of dry THF and transferred to a cooled (0°C) suspension of sodium hydride (1.0 mmol, 0.024 g) in 60 mL of THF. When hydrogen evolution ceased (after 1 hour) trimethylsilyl chloride (1 mmol, 0.109 g) was added by syringe. The mixture became cloudy and was stirred at room temperature for 1 hour before the solvent was removed in vacuo. The remaining residue was partitioned between 10% sodium bicarbonate solution and pentane. The organic layer was washed twice with water and dried over magnesium sulfate, filtered and evaporated to leave a colorless oil (0.204 g, 0.34 mmol, 67%). ¹H NMR (CDCl₃, 360 MHz) 8.68 (broad s, 1 H);

-25-

6.15 (d, 2 H, $J = 2.7$ Hz); 5.13 (dd, 2 H, $J_1 = 18.8$ Hz, $J_2 = 5.0$ Hz); 0.03 (s, 18 H).

Example 13

Preparation of 2-(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsiloxybutyl)pyrrole

A solution of 2-(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole (2.6514 g, 10 mmol) dissolved in 25 ml of dry THF was transferred by canula to a suspension of sodium hydride (0.2640 g, 11 mmol) in 50 ml of THF. The mixture was stirred for 2 hours at room temperature. Trimethylsilyl chloride (1.195 g, 11 mmol) was added dropwise and the solution was stirred overnight. The resulting suspension was evaporated almost to dryness and 50 ml of water was added. The mixture was extracted 3x50 mL with pentane and the organic layers were combined and dried over magnesium sulfate. Upon standing the $MgSO_4$ withdrew a brown impurity from solution leaving a colorless organic layer. The solvent was evaporated leaving a light brown oil (3.128 g). 1H NMR showed that the crude material was 96% product by weight. The overall yield was therefore calculated to be 89%. 1H NMR ($CDCl_3$, 360 MHz) 8.43 (broad s, 1 H); 6.83 (m, 1 H); 6.22 (m, 1 H) 6.17 (q, 1 H, $J = 2.8$ Hz) 5.16 (dd, 1 H, $J_1 = 18.5$ Hz, $J_2 = 5.8$ Hz); 0.03 (s, 9 H).

Example 14

Preparation of 5,10,15,20-Tetrakis(perfluoropropyl)porphyrin from 2,5-Bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole

A solution containing 2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole (0.204 g, 0.34 mmol), pyrrole (0.024 ml, 0.34 mmol) in 70 ml of methylene chloride was transferred to a flask containing anhydrous p-toluenesulfonic acid (50 mg). Aliquots were withdrawn periodically and oxidized by DDQ. TLC on the oxidized aliquots were developed using hexane as the eluant. The reaction was stirred for 12 hours, during which the yield of

-26-

porphyrin appeared to level off. The solution was oxidized with DDQ, neutralized with pyridine, and the solvents were evaporated. The remaining residue was dissolved in hexane (to the extent possible) and chromatographed on silica using
5 hexane as eluant. The first band was collected and evaporated to dryness leaving the title compound (15 mg, 9.1%).

Example 15**A. Preparation of 3,4-dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole**

10 A solution of 2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole (5.0 mmol, 2.32 g) dissolved in 40 ml of dry THF containing 5 ml of pyridine was treated with solid N-bromosuccinimide (NBS; 1.78 g, 10 mmol). The mixture was stirred for 2 hours at room temperature and
15 the solvents were evaporated. The remaining solid was treated with aqueous sodium bicarbonate solution and diethyl ether and separated. An additional extraction was carried out with 50 ml of ether. The organic layers were dried over sodium sulfate and the solvent was evaporated to leave an oily solid.
20 The solid was extracted with cold pentane (3 x 10 ml) leaving behind the succinimide. The pentane extracts were combined and evaporated leaving 3,4-dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole in 89% yield. ¹H NMR (CDCl₃, 360 MHz) 9.33 (broad s, 1 H);
25 5.43 (dd, 2 H, J₁ = 17.5, J₂ = 5.6 Hz); 3.30 (broad s, 2 H).

B. Preparation of 3,4-dichloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole

The procedure of Example 15A is repeated, except that N-chlorosuccinimide (NCS) is used in place of NBS.

C. Preparation of 3,4-diiodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole

30 The procedure of Example 15A is repeated, except that N-iodosuccinimide (NIS) is used in place of NBS.

-27-

Example 16

A. Preparation of 3,4-dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole

5 3,4-Dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole (1 mmol, 0.62 g) was dissolved in 25 ml of dry THF and transferred to a cooled (0°C) suspension of sodium hydride (2.0 mmol, 0.048 g) in 100 mL of THF. When hydrogen evolution ceased (after 2 hours)
10 trimethylsilyl chloride (2 mmol, 0.218 g) was added by syringe. The mixture was stirred at room temperature overnight before the solvent was removed in vacuo. The remaining residue was partitioned between 10% sodium bicarbonate solution and ether and the organic layer was dried
15 over magnesium sulfate, filtered and evaporated to leave a brown oil. The product was purified by chromatography on silica gel using hexane as eluant. The product was the first material off the column (r_f = 0.5) The fractions containing this spot were evaporated to leave a colorless oil which
20 solidified upon cooling under vacuum to leave 3,4-dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole as white needles. (380 mg, 50% yield) ^1H NMR (360 MHz, CDCl_3) 8.84 (broad s, 1 H); 5.30 (dd, 2 H, J_1 = 19.5, J_2 = 1.0); 0.06 (s, 18 H).

25 B. Preparation of 3,4-dichloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole

The procedure of Example 16A is repeated, except that 3,4-dichloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole is used in place of 3,4-bromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole.
30

C. Preparation of 3,4-diiodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole

35 The procedure of Example 16A is repeated, except that 3,4-diiodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-

-28-

hydroxybutyl)pyrrole is used in place of 3,4-bromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole.

Example 17

- A. Preparation of 3-bromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole

The reaction is carried out as in Example 15A except that 1 equivalent of NBS is added instead of 2 equivalents.

- B. Preparation of 3-chloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole

The reaction is carried out as in Example 15B except that 1 equivalent of NCS is added instead of 2 equivalents.

- C. Preparation of 3-iodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole

The reaction is carried out as in Example 15C except that 1 equivalent of NIS is added instead of 2 equivalents.

Example 18

- A. Preparation of 3-bromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole

The reaction is carried out as in Example 16A except that the starting pyrrole is 3-bromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl) pyrrole.

- B. Preparation of 3-chloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole

The reaction is carried out as in Example 16B except that the starting pyrrole is 3-chloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl) pyrrole.

- C. Preparation of 3-iodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole

The reaction is carried out as in Example 16C except that the starting pyrrole is 3-iodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl) pyrrole.

Example 19**A. Preparation of 2,3,12,13-tetrabromo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin**

A solution containing 3,4-dibromo-2,5-bis(2',2',3',-
5 3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole
(0.34 mmol), pyrrole (0.024 ml, 0.34 mmol) in 70 ml of
methylene chloride is transferred to a flask containing
anhydrous p-toluenesulfonic acid (50 mg). Aliquots are
withdrawn periodically and oxidized with DDQ. TLC on the
10 oxidized aliquots are developed using hexane as the eluant.
The reaction is stirred for 12 hours. The solution is
oxidized with DDQ, neutralized with pyridine, and the solvents
are evaporated. The remaining residue is dissolved in hexane
(to the extent possible) and chromatographed on silica using
15 hexane as eluant. The first band is collected and evaporated
to dryness leaving 2,3,12,13-tetrabromo-5,10,15,20-
tetrakis(perfluoropropyl)porphyrin.

B. Preparation of 2,3,12,13-tetrachloro-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

20 The procedure of Example 19A is repeated except that
3,4-dichloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-
trimethylsilyloxybutyl)pyrrole is used in place of 3,4-
dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-
trimethylsilyloxybutyl)pyrrole.

25 C. Preparation of 2,3,12,13-tetraiodo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

The procedure of Example 19A is repeated except that
3,4-diiodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-
trimethylsilyloxybutyl)pyrrole is used in place of 3,4-
30 dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-
trimethylsilyloxybutyl)pyrrole.

Example 20

A. Preparation of 2,12-dibromo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin and 2,13-dibromo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

5 The procedure of Example 19A is repeated except that 3-bromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole is used in place of 3,4-dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole. The two isomers are separated by
10 chromatography on silica gel using pentane as eluant.

B. Preparation of 2,12-dichloro-5,10,15,20-tetrakis(perfluoropropyl)porphyrin and 2,13-dichloro-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

15 The procedure of Example 19B is repeated except that 3-chloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole is used in place of 3,4-dichloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole. The two isomers are separated by chromatography on silica gel using pentane as eluant.

20 C. Preparation of 2,12-diiodo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin and 2,13-diiodo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

The procedure of Example 19C is repeated except that
25 3-iodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole is used in place of 3,4-diiodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole. The two isomers are separated by chromatography on silica gel using pentane as eluant.

Example 21

30 A. Preparation of 2-bromo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

A solution containing 3-bromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole (0.34 mmol), pyrrole (0.048 ml, 0.68 mmol) and 2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole (0.34
35 mmol) in 150 ml of methylene chloride is transferred to a

-31-

flask containing anhydrous p-toluenesulfonic acid (50 mg). Aliquots are withdrawn periodically and oxidized with DDQ. TLC on the oxidized aliquots are developed using hexane as the eluant. The reaction is stirred for 12 hours. The solution
5 is oxidized with DDQ, neutralized with pyridine, and the solvents are evaporated. The remaining residue is dissolved in hexane (to the extent possible) and the product is separated from the unbrominated and dibrominated porphyrin products by chromatography on silica using hexane as eluant.

10 **B. Preparation of 2-chloro-5,10,15,20-tetrakis(perfluoropropyl)porphyrin**

The procedure of Example 21A is repeated except that 3-chloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole is used in place of 3-bromo-
15 2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole.

C. Preparation of 2-iodo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

The procedure of Example 21A is repeated except that
20 3-iodo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole is used in place of 3-bromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole.

Example 22

25 **Preparation of 3,4-disubstituted-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole)**

Palladium catalyzed cross-coupling is performed between 3,4-dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole and a suitably-substituted metal
30 reagent (e.g., R-MgX, R-ZnX, R-borate, R-SnR₃) according to the procedure described by Bray, et al., J. Org. Chem. 1990, 55, 6317-6328.

Example 23

Preparation of 3-substituted-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole)

Palladium catalyzed cross-coupling is performed
5 between 3-bromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole and a suitably-substituted metal reagent (e.g., R-MgX, R-ZnX, R-borate, R-SnR₃) according to the procedure described by Bray, et al., *J. Org. Chem.* 1990, 55, 6317-6328.

10 Example 24

Preparation of [2,3,12,13-tetrakis(perfluorophenyl)-5,10,15,20-tetrakis(perfluoropropyl)porphyrinato]zinc(II)

[2,3,12,13-Tetrabromo-5,10,15,20-tetrakis-
(perfluoropropyl)porphyrinato]zinc(II) (0.2 mmol),
15 pentafluorophenylzinc bromide (10 mmol), and
bis(diphenylphosphino)ferrocene palladium (0.02 mmol) are
dissolved in 10 ml of dry THF and the solution is stirred at
room temperature under nitrogen. Aliquots are withdrawn
periodically and analyzed by TLC on silica gel. Upon
20 completion of the reaction (as indicated by the appearance of
a spot on the TLC plate that fluoresces under long-wavelength
UV light) the solvent is removed and the mixture is
chromatographed on silica gel using hexane as eluant. The
fluorescent band is isolated and the solvent is evaporated to
25 leave [2,3,12,13-tetrakis(perfluorophenyl)-5,10,15,20-
tetrakis(perfluoropropyl)porphyrinato]zinc(II).

Example 25

Preparation of a chiral, nonstrapped porphyrin oxidation catalyst.

30 [2,3,12,13-Tetrabromo-5,10,15,20-tetrakis-
(perfluoropropyl)porphyrinato]zinc(II) (0.2 mmol), 2,6-
dimethoxyphenylzinc chloride (10 mmol), and bis(diphenylphos-
phino)ferrocene palladium (0.02 mmol) are dissolved in 10 ml
of dry THF and the solution is stirred at room temperature
35 under nitrogen. Aliquots are withdrawn periodically and

- 33 -

analyzed by TLC on silica gel. Upon completion of the reaction (as indicated by the appearance of a spot on the TLC plate that fluoresces under long-wavelength UV light) the solvent is removed and the mixture is chromatographed on silica gel using hexane as eluant. The fluorescent band is isolated and the solvent is evaporated to leave [2,3,12,13-tetrakis(2,6-dimethoxyphenyl)-5,10,15,20-tetrakis(perfluoropropyl)porphyrinato]zinc(II). The porphyrin is treated with 0.9 equivalents of boron triiodide in methylene chloride to remove one methyl group (of the eight). The reaction mixture is chromatographed on silica gel using hexane as eluant. The isolated monohydroxy compound (a racemic mixture) is enantiomerically resolved by recrystallization in the presence of a chiral alkaloid (such as quinine) or by chiral stationary phase HPLC.

Example 26

Preparation of an electron deficient face-to-face porphyrin dimer.

[2-Bromo-5,10,15,20-tetrakis(perfluoropropyl)-porphyrinato]zinc(II) (0.2 mmol), 1,8-anthracenedizinc chloride (0.1 mmol), and bis(diphenylphosphino)ferrocene palladium (0.02 mmol) are dissolved in 5 ml of dry THF and the solution is stirred at room temperature under nitrogen. Aliquots are withdrawn periodically and analyzed by TLC on silica gel. Upon completion of the reaction (as indicated by the disappearance of a spots on the TLC plate that corresponds to the starting materials) the solvent is removed and the mixture is chromatographed on silica gel using hexane as eluant. The product band is isolated and the solvent is evaporated to leave 1,8-bis[2'-(5',10',15',20'-tetrakis(perfluoropropyl)porphyrinato(Zn))anthracene.

Example 27**A. Preparation of 2,3,7,8,12,13,17,18-octabromo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin**

3,4-Dibromopyrrole (1mmol) and 3,4-dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole (1 mmol) are dissolved in 200 ml of CH₂Cl₂, and transferred by canula to a flask containing 100 mg of p-toluenesulfonic acid. Aliquots are withdrawn periodically and oxidized with DDQ. TLC on the oxidized aliquots are developed using hexane as the eluant. The reaction is stirred for 12 hours. The solution is oxidized with DDQ, neutralized with pyridine, and the solvents are evaporated. The remaining residue is dissolved in hexane (to the extent possible) and the product is separated from the nonporphyrin products by chromatography on silica using hexane as eluant.

B. Preparation of 2,3,7,8,12,13,17,18-octachloro-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

The procedure of Example 27A is repeated except that 3,4-dichloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole and 3,4-dichloropyrrole are used in place of 3,4-dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole and 3,4-dibromopyrrole.

C. Preparation of 2,3,12,13-Tetrabromo-7,8,17,18-tetrachloro-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

The procedure of Example 27A is repeated except that 3,4-dichloropyrrole is used in place of 3,4-dibromopyrrole.

D. Preparation of 2,3,12,13-Tetraiodo-7,8,17,18-tetrabromo-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

The procedure of Example 27A is repeated except that 3,4-diiodopyrrole is used in place of 3,4-dibromopyrrole.

E. Preparation of 2,3,12,13-Tetraiodo-7,8,17,18-tetrachloro-5,10,15,20-tetrakis(perfluoropropyl)porphyrin

The procedure of Example 27D is repeated except that 3,4-dichloro-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole is used in place of 3,4-

-35-

dibromo-2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsilyloxybutyl)pyrrole.

Example 28**Preparation of 2-(2',2',3',3',4',4',4'-heptafluoro-1'-tosylbutyl)pyrrole**

A solution of 2-(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole (2.6514 g, 10 mmol) dissolved in 25 ml of dry THF is transferred by canula to a suspension of sodium hydride (0.2640 g, 11 mmol) in 50 ml of THF. The mixture is stirred for 2 hours at room temperature. A solution of tosylchloride (11 mmol) in 10 ml of THF is added dropwise and the solution stirred overnight. The resulting suspension is evaporated almost to dryness and 50 ml of water is added. The mixture was extracted 3x50 mL with pentane and the organic layers were combined and dried over magnesium sulfate. The mixture is filtered and the solvent evaporated to yield the product 2-(2',2',3',3',4',4',4'-heptafluoro-1'-tosylbutyl)pyrrole.

Example 29**Preparation of bis-1-(2'-pyrrolyl)-1-perfluoropropylmethane**

A solution of 2-(2',2',3',3',4',4',4'-heptafluoro-1'-tosylbutyl)pyrrole (10 mmol) dissolved in 25 ml of dry THF. Dry pyrrole (40 mmol) is added and the solution is brought to reflux under N₂ for a period of 6 hours. The solution is brought to room temperature with continued stirring and the volatiles removed under vacuum. The solid is extracted with CH₂Cl₂, pumped dry, and chromatographed on silica to give bis-1-(2'-pyrrolyl)-1-perfluoropropylmethane.

Example 30**Preparation of 1,12-perfluoropropyl-1,12-(2'-pyrrolyl)-1,12-tosyl-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.**

The reaction is carried out as in Example 1, except that the starting aldehyde is eicosfluorododecyl-1,12-dial

-36-

dihydrate. 1,12-Perfluoropropyl-1,12-(2'-pyrrolyl)-1,12-hydroxy-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane is isolated as an oil, dissolved in THF and reacted with tosylchloride as in Example 28 to give
5 desired product.

Example 31

Preparation of 1,12-bis(2'-pyrrolyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.

1,12-Perfluoropropyl-1,12-(2'-pyrrolyl)-1,12-tosyl-
10 2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane is reacted with pyrrole as in Example 29. Removal of solvent followed by chromatography on silica gel yields the desired product.

Example 32

15 **Preparation of 1,12-bis[5'-10,15,20-tri(perfluoropropyl)porphyrin]-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.**

1,12-Perfluoropropyl-1,12-(2'-pyrrolyl)-1,12-tosyl-
2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluoro-
20 dodecane is reacted with 2-(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)pyrrole under conditions similar to that described in Example 7. The product, 1,12-bis(5'-10,15,20-perfluoropropylporphyrin)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane is isolated following
25 chromatography on silica gel.

Example 33

Preparation of 1,12-[5',15'-10',20'-di(perfluoropropyl)porphyrin]-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.

30 1,12-Bis(2'-pyrrolyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane is reacted with 2 equivalents of heptafluorobutyraldehyde hydrate and excess sodium hydroxide under conditions similar to Example 3. After a similar work up, the isomeric products are dissolved dry

-37-

benzene and undergo a *p*-toluenesulfonic acid-catalyzed reaction under conditions similar to that described in Example 7. After a work up again similar to that described in Example 7, the product strapped porphyrin obtained from an intramolecular condensation is separated from those products derived from intermolecular reactions by silica gel chromatography.

Example 34

Preparation of 1,12-[5',15'-10',20'-porphyrinyl]-bis(2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane

1,12-Bis(2'-pyrrolyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane is reacted with eicosfluorododecyl-1,12-dial dihydrate under conditions similar to Example 30. As in Example 30, the tetrapyrrolic compound is isolated as the ditosyl derivative and is purified by silica gel chromatography. The material is converted to the product bis-strapped porphyrin and purified utilizing a procedure similar to that described in Example 33.

Example 35

Preparation of a perfluorocarbon-strapped porphyrin with a potential axial ligand for (porphyrinato)metal complexes. (An example of an electron-deficient porphyrin featuring a perfluorocarbon strap linking the porphyrin 5- and 15-(*meso*) positions and a perfluorocarbon strap incorporating an axial ligand linking the porphyrin 10- and 20-positions.)

1,12-Bis(2'-pyrrolyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11 eicosfluorododecane is reacted with 2,5-bis(1',1',2',2',3',-3',4',4'-octafluoropentyl-5'-al)pyridine dihydrate under conditions similar to Example 30. As in Example 30, the tetrapyrrolic compound is isolated as the ditosyl derivative and is purified by silica gel chromatography. The material is converted to the product bis-strapped porphyrin and purified utilizing a procedure similar to that described in Examples 33 and 34.

-38-

Example 36

Preparation of 2-(2",2",3",3",4",4",4"-heptafluoro-1"-hydroxybutyl)dipyrrylmethane.

Dipyrrylmethane is reacted with heptafluoro-
5 butyraldehyde hydrate in a concentrated benzene solution under conditions similar to that described in Example 1. A similar work up followed by silica gel chromatography yields the product 2-(2',2',3',3',4',4',4'-heptafluoro-1'-hydroxybutyl)dipyrrylmethane.

10 **Example 37**

Preparation of 2,2'-bis(2",2",3",3",4",4",4"-heptafluoro-1"-hydroxybutyl)dipyrrylmethane.

Dipyrrylmethane is reacted with heptafluoro-
butyraldehyde hydrate in a concentrated benzene solution under
15 conditions similar to that described in Example 3. A similar work up followed by silica gel chromatography yields the product 2,2'-bis(2",2",3",3",4",4",4"-heptafluoro-1"-hydroxybutyl)dipyrrylmethane.

Example 38

20 Preparation of 5,15-diperfluoropropylporphyrin from 2-(2",2",3",3",4",4",4"-heptafluoro-1"-hydroxybutyl)-dipyrrylmethane.

2-(2",2",3",3",4",4",4"-Heptafluoro-1"-hydroxy-
butyl)dipyrrylmethane is reacted with itself under conditions
25 similar to that described in Example 7. Purification of the product porphyrin is also accomplished in a similar manner as well to give 5,15-diperfluoropropylporphyrin.

Example 39

Preparation of 5,15-diperfluoropropylporphyrin from 2,2'-
30 bis(2",2",3",3",4",4",4"-heptafluoro-1"-hydroxybutyl)-dipyrrylmethane.

2,2'-Bis(2",2",3",3",4",4",4"-heptafluoro-1"-hydroxybutyl)dipyrrylmethane is reacted with dipyrrylmethane under conditions similar to that described in Example 14. A

similar work up and purification gives 5,15-diperfluoropropyl-porphyrin.

Example 40

Preparation of 1,12-(2',2"-dipyrrylmethyl)-2,2,3,3,4,4,5,5,-
5 6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.

Dipyrrylmethane is reacted with eicosfluorododecyl-1,12-dial dihydrate under conditions similar to Example 1, except that the reaction is run in benzene to minimize the oligomeric side products. The product dipyrryl species, 1,12-
10 (2',2"-dipyrrylmethyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,-10,10,11,11-eicosfluorododecane is isolated utilizing silica gel chromatography.

Example 41

Preparation of 1,12-(5',15'-porphyrinyl)-2,2,3,3,4,4,5,5,6,6,-
15 7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.

1,12-(2',2"-Dipyrrylmethyl)-2,2,3,3,4,4,5,5,6,6,-
7,7,8,8,9,9,10,10,11,11-eicosfluorododecane is reacted with dipyrryl methane and subsequently purified as described in Example 39 to give the product 1,12-(5',15'-porphyrinyl)-
20 2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.

Example 42

A. Preparation of 1,12-(5',15'-10,20-dibromoporphyrinyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-
25 eicosfluorododecane.

1,12-(5',15'-Porphyrinyl)-2,2,3,3,4,4,5,5,6,6,7,7,-
8,8,9,9,10,10,11,11-eicosfluorododecane is reacted with N-bromosuccinimide (NBS) in dry CH₂Cl₂ containing excess pyridine. The mixture is stirred at room temperature for two
30 hours and the solvents evaporated. The remaining solid is treated with aqueous sodium bicarbonate solution and CHCl₃, and separated. An additional extraction is carried out with CHCl₃. The organic layers are dried over sodium sulfate, the solvent evaporated and the material is chromatographed on

-40-

silica to enable the separation of the mono- and bis-halogenated porphyrins.

B. Preparation of 1,12-(5',15'-10,20-dichloro-porphyrin)-2,2,3,3,4,4,5,5,6,6,7,7,8,8, 9,9,10,10,11,11-
5 eicosfluorododecane.

The procedure of Example 42A is repeated except that NCS is used in place of NBS.

C. Preparation of 1,12-(5',15'-10,20-diiodo-porphyrin)-2,2,3,3,4,4,5,5,6,6,7,7,8,8, 9,9,10,10,11,11-
10 eicosfluorododecane.

The procedure of Example 42A is repeated except that NIS is used in place of NBS.

Example 43

A. Preparation of 5,15-bis(perfluoropropyl)-10,20-
15 dibromoporphyrin.

5,15-diperfluoropropylporphyrin is reacted with N-bromosuccinimide (NBS) in dry CH_2Cl_2 containing excess pyridine. The mixture is stirred at room temperature for two hours and the solvents evaporated. The remaining solid is
20 treated with aqueous sodium bicarbonate solution and CHCl_3 , and separated. An additional extraction is carried out with CHCl_3 . The organic layers are dried over sodium sulfate, the solvent evaporated and the material is chromatographed on silica to enable the separation of the mono- and bis-
25 halogenated porphyrins.

B. Preparation of 5,15-bis(perfluoropropyl)-10,20-dichloroporphyrin.

The procedure of Example 43A is repeated except that NCS is used in place of NBS.

C. Preparation of 5,15-bis(perfluoropropyl)-10,20-diiodoporphyrin.
30

The procedure of Example 43A is repeated except that NIS is used in place of NBS.

Example 44

Preparation of 2,3-bis(perfluoromethyl)pyrrole from 1-triisopropylsilyl-2,3-dibromopyrrole.

1-Triisopropylsilyl-2,3-dibromopyrrole is reacted
5 with a twenty-fold molar excess of trifluoromethylzinc chloride in the presence of 5 mole percent Pd(PPh₃)₄ in tetraglyme at 100°C. The reaction is monitored by TLC. After 48 hours, the reaction is quenched with water, the solvents removed, and the material is chromatographed on silica to
10 yield 1-triisopropylsilyl-2,3-diperfluoromethylpyrrole. This material is desilylated with tetrabutylammonium fluoride and then used immediately in a porphyrin condensation reaction.

Example 45

Preparation of 1,6-(3',4'-pyrrolyl)-1,1,2,2,3,3,4,4,5,5,6,6-
15 dodecafluorohexane.

Similar to that described in Example 44, a dilute solution of 1-triisopropylsilyl-2,3-dibromopyrrole is reacted with one equivalent of 1,6-bis(chlorozincyl)-1,1,2,2,3,3,4,4,5,5,6,6-dodecafluorohexane in tetraglyme in the presence
20 of Pd catalyst for one week at 100°C. Work up is as described in Example 44 above.

Example 46

Preparation of 1,6-(3',4'-pyrrolyl)-1,1,2,2,3,3,4,4,5,5,6,6-dodecafluorohexane from pyrrole and perfluorocyclooctyl-4-yne.

25 Pyrrole and perfluorocyclooctyl-4-yne are reacted in CHCl₃ under conditions appropriate to induce a Diels-Alder reaction between the two reactants that is followed by a reverse Diels-Alder reaction that extrudes acetylene.

Example 47

30 Preparation of 1,6-[2'-(2'',2'',3'',3'',4'',4'',4''-heptafluoro-1''-hydroxybutyl)-3',4'-pyrrolyl]-1,1,2,2,3,3,4,4,5,5,6,6-dodecafluorohexane

1,6-(3',4'-Pyrrolyl)-1,1,2,2,3,3,4,4,5,5,6,6-octafluorohexane is reacted with heptafluorobutyraldehyde.

-42-

hydrate according to the procedure outlined in Example 1; purification and isolation are accomplished according to the general scheme outlined in Example 1 as well.

Example 48

- 5 Preparation of 2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsiloxybutyl)-3-trifluoromethylpyrrole.

Trifluoromethylzinc chloride is reacted with 3-bromo-2,5-bis(2,2,3,3,4,4,4-heptafluoro-1-trimethylsilyloxybutyl)pyrrole utilizing reaction and
10 purification conditions described in Example 22.

Example 49

Preparation of 2-(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsiloxybutyl)-3-trifluoromethylpyrrole.

2-(2',2',3',3',4',4',4'-Heptafluoro-1'-
15 trimethylsiloxybutyl)pyrrole is reacted with N-trifluoromethyl-N-nitrosotrifluoromethanesulfonamide under conditions similar to those described by Umemoto, et al., *Bull. Chem. Soc. Jpn.* 1986, 59, 447. Following the recommended work up procedure, the solvent is removed and the
20 organic residue chromatographed on silica to yield 2-(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsiloxybutyl)-3-trifluoromethylpyrrole.

Example 50

- Preparation of 1,12-(2',5'-Pyrrolyl)-1,12-trimethylsiloxy-
25 2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10-eicosofluorodecane from eicosfluorododecyl-1,12-dial dihydrate.

Pyrrole is reacted with one equivalent of eicosfluorododecyl-1,12-dial dihydrate under conditions similar to that described in Example 40. Following a similar
30 work up procedure, silica gel chromatography enables the separation of the product 1",1"-[2',5'-bis(2",2",3",3",4",4",4"-heptafluoro-1"-trimethylsiloxybutyl)pyrrolyl]-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane from 1,12-perfluoropropyl-1,12-(2'-pyrrolyl)-1,12-hydroxy-

-43-

2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane and other oligomeric by products. The product is converted to its trimethylsiloxy derivative utilizing an experimental procedure similar to that described in Example 12.

Example 51

Preparation of 1,12-(2',5'-pyrrolyl)-1,12-trimethylsiloxy-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10-eicosfluorododecane from eicosfluorododecyl-1,12-diacid chloride.

10 Pyrrole is reacted with one equivalent of eicosfluorododecyl-1,12-diacid chloride in dry THF under nitrogen. The reaction is monitored by TLC. At the reaction endpoint, the solvents are removed and the material chromatographed on silica to give the fluorocarbon-strapped 2,5-dialdehydyl pyrrole. The product is reacted with sodium borohydride in THF, quenched with water and re-chromatographed on silica, isolated, and converted to its trimethylsiloxy derivative utilizing an experimental procedure similar to that described in Example 12 to give the desired product, 1,12-[2',5'-bis(1"-2",2",3",3",4",4",4"-heptafluoro-1"-trimethylsiloxybutyl)pyrrolyl]-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane. A large number of pyrrolic precursors suitable for the preparation of perfluorocarbon-strapped porphyrins exist using the condensation methodologies described herein. A number of representative pyrrolyl and dipyrrolyl species are described in Examples 53-55.

Example 52

Preparation of cis- and trans-1,12-[5',10'-15',20'-porphyrinyl]-bis(2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane

1,12-[2',5'-Bis(1"-2",2",3",3",4",4",4"-heptafluoro-1"-trimethylsiloxybutyl)pyrrolyl]-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane is reacted with pyrrole under conditions similar to that described in Example 14. At the reaction endpoint, work up follows the procedure

-44-

outlined in Example 14; the two product porphyrins, cis- and trans-1,12-[5',10'-15',20'-porphyrinyl]-bis(2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane) are separated and isolated by chromatography on silica.

5 Example 53

Preparation of 1,12-bis(3'-pyrrolyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane from perfluorocarbonbis(chlorozinc) reagents.

Excess 3-bromopyrrole is reacted with 1,12-bis(chlorozincyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane under conditions similar to that described in Example 45. Work up and chromatographic isolation affords 1,12-bis(3'-pyrrolyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.

15 Example 54

Preparation of 1,12-bis[3'-2'-(2'',2'',3'',3'',4'',4'',4''-heptafluoro-1''-trimethylsiloxybutyl)pyrrolyl]-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane from perfluorocarbonbis(chlorozinc) reagents.

4-Bromo-2-(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsiloxybutyl)pyrrole is reacted with 1,12-bis(chlorozincyl)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane under conditions similar to that described in Example 45. Work up and chromatographic isolation affords 1,12-bis[3'-2'-(2'',2'',3'',3'',4'',4'',4''-heptafluoro-1''-trimethylsiloxybutyl)pyrrolyl]-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.

Example 55

Preparation of 1,12-bis[3'-2'-(2'',2'',3'',3'',4'',4'',4''-heptafluoro-1''-trimethylsiloxybutyl)pyrrolyl]-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane from 1,12-bis(N'-N-nitrosotrifluoromethanesulfonamide)-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane.

-45-

Excess 2-(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsiloxybutyl)pyrrole is reacted with 1,12-bis(N'-N-nitrosotrifluoromethanesulfonamide)-2,2,3,3,4,4,5,5,6,6,-7,7,8,8,9,9,10,10,11,11-eicosfluorododecane under conditions similar to that described for Example 49. Work up and chromatographic separation yields 1,12-bis[3'-2'-(2'',2'',-3'',3'',4'',4'',4''-heptafluoro-1''-trimethylsiloxybutyl)pyrrolyl]-2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11-eicosfluorododecane. Straps originating and/or terminating at a pyrrole 3- or 4- position will undoubtedly give rise to a large number of isomers when converted to their 2-(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsiloxybutyl) derivatives. Cyclization of these materials, the 2,5-bis(2',2',3',3',4',4',4'-heptafluoro-1'-trimethylsiloxybutyl) derivatives or simple perfluorocarbon-linked pyrroles (Example 53) will allow for the formation of mono-strapped porphyrins along with cis and trans-strapped species. A variety of isomeric derivatives of these materials will be present prior to chromatographic separation. One particular advantage of strapping porphyrins at their β -positions is that it leaves the *meso* positions free for further derivatization.

Example 56

Preparation Of Chiral, Nonstrapped Porphyrin Oxidation Catalyst.

2,3,12,13-Tetrabromo-5,10,15,20-tetrakis-(perfluoropropyl)porphyrinato(zinc) (0.2 mmol), 2,6-dimethoxyphenylzinc chloride (10 mmol), and bis(diphenylphosphino)ferrocene palladium (0.02 mmol) are dissolved in 10 ml of dry THF and the solution is stirred at room temperature under nitrogen. Aliquots are withdrawn periodically and analyzed by TLC on silica gel. When a substantial portion of the material has three bromine atoms exchanged for three aryl moieties, the reaction is quenched by the introduction of air. The solvent is removed and the mixture is chromatographed on silica gel using hexane THF as eluant. The appropriate band is isolated and the solvent is

-46-

evaporated to leave 2,3,12,-tris(2,6 dimethoxyphenyl)-13-bromo-5,10,15,20-tetrakis(perfluoropropyl)porphyrinato(zinc). The porphyrin is treated with 2-t-butylphenylzinc chloride (10 mmol), and bis(diphenylphosphino)ferrocene palladium (0.02 mmol) dissolved in 10 ml of dry THF and the solution is stirred at room temperature under nitrogen. The reaction mixture is chromatographed on silica gel using hexane as eluant. The isolated a racemic mixture is enantiomerically resolved by recrystallization in the presence of a chiral alkaloid (such as quinine) or by chiral stationary phase HPLC.

Example 57

Preparation of Nonstrapped- And Strapped-porphyrin Oxidation Catalysts

The isolation of chiral catalysts relies on the demonstrated enhanced solubility imparted by the perfluoroalkyl groups. Porphyrins (such as tetraphenylporphyrin (TPP)) are notoriously difficult to chromatograph. Perfluoroalkyl-substituted porphyrins, on the other hand, are easily purified by normal phase or reverse phase HPLC. Therefore, a wide variety of commercially available chiral chromatography columns are potentially applicable to these compounds, along with their more highly-elaborated derivatives. This capability simplifies the general preparative scheme since racemic mixtures of the catalysts may be prepared and resolved yielding catalysts with opposite stereoselectivity. Two general approaches to asymmetric catalyst synthesis are outlined in Figures 1 and 2, wherein: M = Zn or H₂; X = Br or I; R = H, O-methyl, O-ethyl, O-isopropyl, Cl, methyl, ethyl, isopropyl, or tert-butyl; Y = ZnCl or B(OH)₂; R_F = perfluoroalkyl; R' = a group different from R; L₂ = a phosphorous-, nitrogen- or arsenic-containing Lewis base; and m = 4-12. In the first method, shown in Figure 1, three identical bulky groups are appended to the porphyrin, and the resulting singly halogenated species is isolated. Chirality is imparted by the attachment of the last, unsymmetrically substituted aryl group that also

-47-

contains an axial ligand tethered to the 2-position. The axial ligand will effectively close off one face of the molecule for epoxidation (or hydroxylation) chemistry. The resulting racemic product can be resolved by chiral HPLC.

5 Resolution can be further enhanced by the addition of chiral amine bases that can act as axial ligands to form diastereomeric complexes for achiral reverse phase media. The second general approach to enantioselective epoxidation (or chiral alkane hydroxylation) catalysts involves the

10 preparation of strapped porphyrins, as outlined in Figure 2. The advantages to this scheme are that entry to the metal center is extremely restricted and that fluorocarbon containing straps may be used. (The incorporation of chemically inert perfluoroalkyl straps is key if

15 intramolecular reactions are implicated in catalyst degradation.) Molecular models indicate that the nine and ten atom bridges can be accommodated without severely distorting the porphyrin core. Coupling of the final two groups and resolution of the enantiomers will be carried out as above.

20 The effectiveness of the catalysts for enantioselective epoxidations (or selective alkane hydroxylations) can be evaluated by chiral HPLC, optical rotation, and shift reagent NMR techniques. Enantioselectivity as well as absolute product chirality can be monitored as a function of the

25 turnover number to determine if partially degraded, catalytically competent species are formed.

Those skilled in the art will appreciate that numerous changes and modifications may be made to the preferred embodiments of the invention and that such changes

30 and modifications may be made without departing from the spirit of the invention. For example, it is believed that the methods of the present invention can be practiced using porphyrin-related compounds such as chlorins, phorbins, bacteriochlorins, porphyrinogens, sapphyrins, texaphrins, and

35 pthalocyanines in place of porphyrins. It is therefore intended that the appended claims cover all such equivalent


-48-

variations as fall within the true spirit and scope of the invention.


-49-

WHAT IS CLAIMED IS:

1. A synthetic process comprising contacting a pyrrole derivative having formula:



with a compound having formula:



in organic solvent for a time and under conditions effective to form a covalent linkage between said pyrrole derivative and said pyrrole, wherein:

each L is a leaving group;

n is 0, 1, or 2; and

R_A and R_B are defined such that:

at least one R_A is a group that is electron-withdrawing relative to hydrogen;

at least two R_A, together, form a covalently-bound, multi-atom linkage;

at least two R_B, together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B, together, form a covalently-bound, multi-atom linkage.

2. The process of claim 1 wherein R_A is alkyl having 1 to about 50 carbon atoms, aryl having 3 to about 75 carbon atoms, CF₃R₁, N-(alkyl)₃, NH₃⁺, NO₂, SO₂-(alkyl), CN, SO₂-(aryl), C(O)OH, F, Cl, Br, I, C(O)O-(alkyl), C(O)-(alkyl), or CHO, wherein R₁ is alkyl, alkenyl, alkynyl, or a halogenated derivative thereof, alkyl groups have from about

-50-

1-50 carbon atoms, alkenyl and alkynyl groups have from about 2-50 carbon atoms, and aryl groups have about 3-75 carbon atoms.

3. The process of claim 1 wherein R_A is haloalkyl having 1 to about 20 carbon atoms or haloaryl having 6 to about 20 carbon atoms.

4. The process of claim 1 wherein R_A is perfluoroalkyl having 5 to about 20 carbon atoms or perfluoroaryl having 6 to about 20 carbon atoms.

5. The process of claim 1 wherein at least two R_A , together, form a covalently-bound, multi-atom linkage.

6. The process of claim 5 wherein at least two R_A , together, are alkyl having 3 to about 35 carbon atoms, aryl having 6 to about 40 carbon atoms, $C(O)O-(alkyl)$ or $C(O)-(alkyl)$ wherein alkyl groups have from about 1-50 carbon atoms and aryl groups have about 3-75 carbon atoms.

7. The process of claim 5 wherein at least two R_A , together, are haloalkyl having 3 to about 35 carbon atoms or haloaryl having 6 to about 40 carbon atoms.

8. The process of claim 5 wherein at least two R_A , together, are perfluoroalkyl having 3 to about 35 carbon atoms or perfluoroaryl having 6 to about 50 carbon atoms.

9. The process of claim 1 wherein at least two R_B , together, form a covalently-bound, multi-atom linkage.

10. The process of claim 9 wherein at least two R_B , together, are alkyl having 3 to about 35 carbon atoms, aryl having 6 to about 40 carbon atoms, $C(O)O-(alkyl)$ or $C(O)-(alkyl)$ wherein alkyl groups have from about 1-50 carbon atoms and aryl groups have about 3-75 carbon atoms.

-51-

11. The process of claim 9 wherein at least two R_B , together, are haloalkyl having 3 to about 35 carbon atoms or haloaryl having 6 to about 40 carbon atoms.


12. The process of claim 9 wherein at least two R_B , together, are perfluoroalkyl having 3 to about 35 carbon atoms or perfluoroaryl having 6 to about 50 carbon atoms.

13. The process of claim 1 wherein at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

14. The process of claim 1 wherein L is a halogen, a sulfate, a sulfonate, or a carboxylate.

15. The process of claim 1 further comprising oxidizing said reaction mixture.

16. The process of claim 1 further comprising isolating from said reaction mixture a composition containing a porphyrinogen having formula:




17. The process of claim 16 further comprising oxidizing said porphyrinogen for a time and under reaction conditions effective to form a corresponding porphyrin.

18. The product of the process of claim 1.

-52-

19. The product of the process of claim 16.
20. The product of the process of claim 17.
21. A synthetic process comprising reacting at least two molecules having formula:



in organic solvent for a time and under conditions effective to form a covalent bond between said molecules, wherein:

L is a leaving group;

n is 0, 1, or 2; and

R_A and R_B are defined such that:


at least one R_A is a group that is electron-withdrawing relative to hydrogen;

at least two R_A, together, form a covalently-bound, multi-atom linkage;

at least two R_B, together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B, together, form a covalently-bound, multi-atom linkage.

22. A synthetic process comprising the steps of:
- (a) contacting an aldehyde having formula R_A-CHO with a pyrrole derivative having formula:



in organic solvent in the presence of acid for a time and under conditions effective to form a reaction mixture

-53-

comprising water and an adduct of said aldehyde and said pyrrole derivative; and

(b) removing from said reaction mixture a portion of said water;

wherein q is 0, 1, or 2 and:

at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

23. The process of claim 22 wherein at least two R_A , together, form a covalently-bound, multi-atom linkage.

24. The process of claim 22 wherein at least two R_B , together, form a covalently-bound, multi-atom linkage.

25. The process of claim 22 wherein at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

26. The process of claim 22 wherein said solvent is nonpolar and aprotic.

27. The process of claim 22 wherein said solvent forms an azeotrope with water.

28. The process of claim 22 wherein said solvent is benzene, toluene, xylenes, methylene chloride, chloroform, trichloroethylene, or a mixture thereof.

29. The process of claim 22 wherein said acid is a non-volatile, protic acid.

30. The process of claim 22 wherein said acid is benzoic acid, *p*-toluenesulfonic acid, methanesulfonic acid,

-54-

trifluoroacetic acid, boron trifluoride, boron trichloride, or a mixture thereof.

31. The process of claim 22 wherein a catalytic amount of said acid is used.

32. The process of claim 27 wherein said water is removed by distilling said azeotrope.

33. The process of claim 32 further comprising collecting said distilled azeotrope in a vessel such that said distilled azeotrope forms a substantially aqueous phase and a substantially organic phase.

34. The process of claim 33 further comprising returning said substantially organic phase to said reaction mixture.

35. The process of claim 31 further comprising contacting said distilled azeotrope with a drying agent for a time sufficient to remove a portion of said water contained therein.

36. The process of claim 35 further comprising returning said dried distillate to said reaction mixture.


37. The process of claim 35 wherein said drying agent comprises phosphorous pentoxide, calcium hydride, calcium oxide, barium oxide, lithium aluminum hydride, molecular sieves, or a mixture thereof.

38. The process of claim 32 further comprising adding organic solvent to said reaction mixture.

39. The process of claim 22 further comprising oxidizing said reaction mixture.


-55-

40. The process of claim 22 further comprising isolating from said reaction mixture a composition containing a porphyrinogen having formula:



41. The process of claim 40 further comprising oxidizing said porphyrinogen for a time and under reaction conditions effective to form a corresponding porphyrin.

42. A synthetic process comprising the steps of:
 (a) reacting an alcohol having formula:



in organic solvent in the presence of acid for a time and under conditions effective to form a reaction mixture comprising water and an adduct of said alcohol; and

(b) removing from said reaction mixture a portion of said water;

wherein n is 0, 1, or 2 and:


at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or


at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

-56-

43. A synthetic process comprising the steps of:
 (a) contacting a pyrrole derivative having formula:



with a compound having formula:



in organic solvent in the presence of acid for a time and under conditions effective to form a reaction mixture comprising water and an adduct of said pyrrole derivative and said compound; and

(b) removing from said reaction mixture a portion of said water;


wherein n is 0, 1, or 2 and:

at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

44. A compound having formula:



wherein:

each L is a leaving group;

n is 0, 1, or 2; and

-57-

R_A and R_B are defined such that:

at least one R_A is a group that is electron-withdrawing relative to hydrogen;

at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

45. The compound of claim 44 wherein R_A is alkyl having 1 to about 50 carbon atoms, aryl having 3 to about 75 carbon atoms, CF_2R_1 , $N-(alkyl)_3$, NH_3 , NO_2 , $SO_2-(alkyl)$, CN , $SO_2-(aryl)$, $C(O)OH$, F , Cl , Br , I , $C(O)O-(alkyl)$, $C(O)-(alkyl)$, or CHO , wherein R_1 is alkyl, alkenyl, alkynyl, or a halogenated derivative thereof, alkyl groups have from about 1-50 carbon atoms, alkenyl and alkynyl groups have from about 2-50 carbon atoms, and aryl groups have about 3-75 carbon atoms.

46. The compound of claim 44 wherein R_A is haloalkyl having 1 to about 20 carbon atoms or haloaryl having 6 to about 20 carbon atoms.

47. The compound of claim 44 wherein R_A is perfluoroalkyl having 5 to about 20 carbon atoms or perfluoroaryl having 6 to about 20 carbon atoms.

48. The compound of claim 44 wherein at least two R_A , together, form a covalently-bound, multi-atom linkage.

49. The compound of claim 48 wherein at least two R_A , together, are alkyl having 3 to about 35 carbon atoms, aryl having 6 to about 40 carbon atoms, $C(O)O-(alkyl)$ or $C(O)-(alkyl)$ wherein alkyl groups have from about 1-50 carbon atoms and aryl groups have about 3-75 carbon atoms.

-58-

50. The compound of claim 48 wherein at least two R_A , together, are haloalkyl having 3 to about 35 carbon atoms or haloaryl having 6 to about 40 carbon atoms.

51. The compound of claim 48 wherein at least two R_A , together, are perfluoroalkyl having 3 to about 35 carbon atoms or perfluoroaryl having 6 to about 50 carbon atoms.

52. The compound of claim 44 wherein at least two R_B , together, form a covalently-bound, multi-atom linkage.

53. The compound of claim 52 wherein at least two R_B , together, are alkyl having 3 to about 35 carbon atoms, aryl having 6 to about 40 carbon atoms, $C(O)O$ -(alkyl) or $C(O)$ -(alkyl) wherein alkyl groups have from about 1-50 carbon atoms and aryl groups have about 3-75 carbon atoms.

54. The compound of claim 52 wherein at least two R_B , together, are haloalkyl having 3 to about 35 carbon atoms or haloaryl having 6 to about 40 carbon atoms.


55. The compound of claim 52 wherein at least two R_B , together, are perfluoroalkyl having 3 to about 35 carbon atoms or perfluoroaryl having 6 to about 50 carbon atoms.

56. The compound of claim 44 wherein at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

57. The compound of claim 44 wherein L is a halogen, a sulfate, a sulfonate, or a carboxylate.

-59-

58. A compound having formula:



wherein:

L is a leaving group;

n is 0, 1, or 2; and

R_A and R_B are defined such that:


at least one R_A is a group that is electron-withdrawing relative to hydrogen;

at least two R_A, together, form a covalently-bound, multi-atom linkage;

at least two R_B, together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B, together, form a covalently-bound, multi-atom linkage.

59. A compound having formula:



wherein q is 0, 1, or 2 and:


at least two R_A, together, form a covalently-bound, multi-atom linkage;

at least two R_B, together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B, together, form a covalently-bound, multi-atom linkage.

- 60 -

60. A compound having formula:




wherein n is 0, 1, or 2 and:

at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

61. A compound having formula:



wherein n is 0, 1, or 2 and:


at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or


at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

- 61 -


62. A compound having formula (7), (8), or (9):



(7)



(8)



(9)

wherein M and M' are metal atoms and:




at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

- 62 -

63. A compound having formula (7), (8), or (9):



wherein M and M' are metal atoms and:

a single R_B on each pyrrole moiety is a halogen; or
both R_B on each of two non-adjacent pyrrole moieties are halogens.

64. In a process in which an alkane is selectively oxidized by contact with air or oxygen in the presence of a catalyst comprising a transition metal coordination complex, the improvement wherein said complex comprises a compound according to claim 62.

65. In a process in which a carbon-carbon double bond is selectively oxidized by contact with air or oxygen in the presence of a catalyst comprising a transition metal

-63-

coordination complex, the improvement wherein said complex comprises a compound according to claim 62.

66. A supported catalyst comprising a compound according to claim 62 on a solid support material.




Figure 1

2/2




Figure 2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/15899

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07D 487/22; C07B 61/00; C07C 29/50, 31/12

US CL : 540/145, 472; 548/400; 568/399, 910, 910.5; 514/185

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 540/145, 472; 548/400; 568/910, 910.5; 514/183, 185

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y, P	US, A, 5,405,957 (TANG et al.) 11 April 1995, see entire document.	1-66
Y, P	JP, A, 6-128263 (YUSUKE IZUMI) 10 May 1994, see Abstract translation.	1-63
Y	JOURNAL OF CHEMICAL SOCIETY, PERKIN-TRANSACTION I; issued April 1974, JACKSON ET AL., "PYRROLES AND RELATED COMPOUNDS PART XXV. PEMPTOPORPHYRIN, ISOPEMPTOPORPHYRIN AND CHLOROCRUOROPORPHYRIN (SPIROGRAPHIS PORPHYRIN)", pages 480-490, see entire document.	1-63
X	US, A, 5,118,886 (ELLIS, JR. et al.) 02 June 1992, see entire document.	40, 41, 62-66



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	* T* later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be part of particular relevance	* X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* &* document member of the same patent family
* O* document referring to an oral disclosure, use, exhibition or other means	
* P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

25 MARCH 1996

Date of mailing of the international search report

13 MAY 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

P.R. SRIPADA aco

Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/15899

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US, A, 5,212,300 (ELLIS, JR. et al.) 18 May 1993, see entire document.	40, 41, 62-66

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/15899

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

I. Claims 1-61, drawn to processes of making pyrrole multimers, classified in 548/400.

II. Claims 62 and 63, drawn to porphyrins classified in 540/145, 472.

III. Claims 64-66, drawn to process as catalyst 568/910.5; 502/315.

The groupings lack unity of invention because Grs. I and II are drawn to pyrroles and porphyrin cores and are classified in different classes and require different searches. Besides, these are non-art recognized equivalents. Reference of Gr. III will not be reference to Gr. I or II; 37 1.475(d).



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 487/22, C07B 61/00, C07C 29/50, 31/12	A1	(11) International Publication Number: WO 96/17844 (43) International Publication Date: 13 June 1996 (13.06.96)
(21) International Application Number: PCT/US95/15899 (22) International Filing Date: 7 December 1995 (07.12.95) (30) Priority Data: 08/352,555 8 December 1994 (08.12.94) US (71) Applicant: THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; Center for Technology Transfer, Suite 300, 3700 Market Street, Philadelphia, PA 19104-3147 (US). (71)(72) Applicant and Inventor: DIMAGNO, Stephen [US/US]; 2828 Jackson Drive, Lincoln, NE 68502 (US). (72) Inventor: THERIEN, Michael, J.; 2411 D Delancey Place, Philadelphia, PA 19103 (US). (74) Agents: ELDERKIN, Dianne, B. et al.; Woodcock Washburn Kurtz MacKiewicz & Norris, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>With amended claims and statement.</i> Date of publication of the amended claims and statement: 22 August 1996 (22.08.96)
(54) Title: NOVEL PORPHYRINS AND PORPHYRIN SYNTHESIS TECHNIQUES (57) Abstract Strapped porphyrins and electron-deficient porphyrins are provided, as well as processes and intermediates for their preparation. In certain embodiments, such compounds are prepared by nucleophilic displacement of leaving groups from methylpyrroles. In other embodiments, such compounds are prepared by condensing pyrrole derivatives and removing water thus formed from the resulting reaction mixture.		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

AMENDED CLAIMS

[received by the International Bureau on 8 July 1996 (08.07.96);
original claims 58 and 60 amended; remaining claims unchanged (3 pages)]

50. The compound of claim 48 wherein at least two R_A , together, are haloalkyl having 3 to about 35 carbon atoms or haloaryl having 6 to about 40 carbon atoms.
51. The compound of claim 48 wherein at least two R_A , together, are perfluoroalkyl having 3 to about 35 carbon atoms or perfluoroaryl having 6 to about 50 carbon atoms.
52. The compound of claim 44 wherein at least two R_B , together, form a covalently-bound, multi-atom linkage.
53. The compound of claim 52 wherein at least two R_B , together, are alkyl having 3 to about 35 carbon atoms, aryl having 6 to about 40 carbon atoms, $C(O)O-(alkyl)$ or $C(O)-(alkyl)$ wherein alkyl groups have from about 1-50 carbon atoms and aryl groups have about 3-75 carbon atoms.
54. The compound of claim 52 wherein at least two R_B , together, are haloalkyl having 3 to about 35 carbon atoms or haloaryl having 6 to about 40 carbon atoms.
55. The compound of claim 52 wherein at least two R_B , together, are perfluoroalkyl having 3 to about 35 carbon atoms or perfluoroaryl having 6 to about 50 carbon atoms.
56. The compound of claim 44 wherein at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.
57. The compound of claim 44 wherein L is a halogen, a sulfate, a sulfonate, or a carboxylate.
58. A compound having formula:



AMENDED SHEET (ARTICLE 19)

wherein:

L is a leaving group;

n is 1 or 2; and

R_A and R_B are defined such that:


at least one R_A is a group that is electron-withdrawing relative to hydrogen;

at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

59. A compound having formula:




wherein q is 0, 1, or 2 and:

at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

60. A compound having formula:




wherein n is 1 or 2; and

at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

61. A compound having formula:





wherein n is 0, 1, or 2 and:

at least two R_A , together, form a covalently-bound, multi-atom linkage;

at least two R_B , together, form a covalently-bound, multi-atom linkage; or

at least one R_A and at least one R_B , together, form a covalently-bound, multi-atom linkage.

62. A compound having formula (7), (8), or (9):



AMENDED SHEET (ARTICLE 19)

STATEMENT UNDER ARTICLE 19

Claims 58 and 60 have been amended. No new matter has been added.





Figure 1

2/2



Continue as in Figure 1

Figure 2

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.